arXiv:hep-th/9506087v1 13 Jun 1995

June 1995
SNUCTP 95 - 66
hep-th/9506087

Duality in Complex sine-Gordon Theory

Q-Han Park[1]

and

H. J. Shin[2]

Department of Physics
and
Research Institute of Basic Sciences
Kyunghee University
Seoul, 130-701, Korea

ABSTRACT

New aspects of the complex sine-Gordon theory are addressed through the reformulation of the theory in terms of the gauged Wess-Zumino-Witten action. A dual transformation between the theory for the coupling constant $\beta > 0$ and the theory for $\beta < 0$ is given which agrees with the Krammers-Wannier duality in the context of perturbed conformal field theory. The Bäcklund transform and the nonlinear superposition rule for the complex sine-Gordon theory are presented and from which, exact solutions, solitons and breathers with U(1) charge, are derived. We clarify topological and nontopological nature of neutral and charged solitons respectively, and discuss about the duality between the vector and the axial U(1) charges.

---

[1] E-mail address; qpark@nms.kyunghee.ac.kr
[2] E-mail address; hjshin@nms.kyunghee.ac.kr

The complex sine-Gordon theory, which generalizes the well-known sine-Gordon theory with an internal U(1) degree of freedom, first appeared as a model of relativistic vortices in a superfluid [1], and also independently in a treatment of O(4) nonlinear sigma model[2]. Recently, Bakas has shown that the complex sine-Gordon theory may be reformulated in terms of the gauged Wess-Zumino-Witten(WZW) action and interpreted the theory as the integrably deformed SU(2)/U(1)-coset model for the parafermions in the large N limit for the level N[3]. This led to subsequent generalizations of the sine-Gordon and the complex sine-Gordon theories to other coset cases[4][5][6] as well as generalizations of solitons and breathers with internal degrees of freedom[7]. Reformulation of the theory as the deformed coset model also provides a natural explanation for the behavior of exact factorizable S-matrix[8].

In this Letter, we report new aspects of the complex sine-Gordon theory which arises from the reformulation of the theory in terms of the gauged WZW action and a specific choice of the gauge described later. We find the exact duality between the theories for the coupling constant $\beta > 0$ and $\beta < 0$. We show that this agrees precisely with the Krammers-Wannier duality between the spin variables s_j and the dual spin variables μ_j of Z_N-parafermion theory[9].[3] We derive the Bäcklund transform and the nonlinear superposition rule for the complex sine-Gordon theory from the gauged WZW action in the gauge $A = \bar{A} = 0$, from which exact solutions, solitons and breathers with U(1) charge, are obtained. We clarify the topological nature of soliton solutions. It is shown that charged solitons are in general non-topological solitons and become topological only when they become neutral. We also address the duality between the axial and the vector U(1) charges of the complex sine-Gordon theory.

The complex sine-Gordon theory in terms of the gauged WZW action is given by

$$\begin{aligned} I(g, A, \bar{A}, \beta) &= I_{WZW}(g) + \frac{1}{2\pi}\int \mathrm{Tr}(-A\bar{\partial}gg^{-1} + \bar{A}g^{-1}\partial g + Ag\bar{A}g^{-1} - A\bar{A}) \\ &- \frac{\beta}{2\pi}\int \mathrm{Tr}gTg^{-1}\bar{T} \end{aligned} \qquad (1)$$

where $I_{WZW}(g)$ is the SU(2)-WZW action for a map $g : M \to \mathrm{SU(2)}$ on two-dimensional Minkowski space M. The connection $A, \bar{A}$ gauge the anomaly free diagonal subgroup U(1) of SU(2) and $T = -\bar{T} = i\sigma_3 = \mathrm{diag}(i, -i)$ where $\sigma_i; i = 1, 2, 3$ are Pauli matrices. This action possesses the local U(1) vector symmetry as well as the global U(1) axial symmetry. The equation of motion of the action (1) takes a zero curvature form,

$$\delta_g I = -\frac{1}{2\pi}\int \mathrm{Tr}[\ \partial + g^{-1}\partial g + g^{-1}Ag + \beta\lambda T\ ,\ \bar{\partial} + \bar{A} + \frac{1}{\lambda}g^{-1}\bar{T}g\ ]g^{-1}\delta g = 0 \qquad (2)$$

which, together with the constraint equation

$$\delta_A I(g, A, \bar{A}) = \frac{1}{2\pi}\int \mathrm{Tr}(\ -\bar{\partial}gg^{-1} + g\bar{A}g^{-1} - \bar{A}\ )\delta A = 0$$

---

[3] The changing sign of coupling constants β under the Krammers-Wannier duality has been pointed out by Bakas and the duality between the two theories was suggested but without an explicit duality transform rule[3].

$$\delta_{\bar{A}} I(g, A, \bar{A}) \;=\; \frac{1}{2\pi}\int \mathrm{Tr}(\; g^{-1}\partial g + g^{-1} A g - A\;)\delta\bar{A} = 0\;, \tag{3}$$

defines the theory at the classical level. In particular, the diagonal part of the equation of motion (2), when combined with the constraint equation, results in the flatness condition of the connection $A, \bar{A}$

$$\partial\bar{A} - \bar{\partial}A = 0. \tag{4}$$

In order to show that Eqs.(2) and (3) are indeed the complex sine-Gordon equation, we introduce a local parametrization of g,

$$g = \begin{pmatrix} u & i\sqrt{1-uu^*}e^{i\theta} \\ i\sqrt{1-uu^*}e^{-i\theta} & u^* \end{pmatrix}\;, \tag{5}$$

and denote that $A = a\sigma_3,\ \bar{A} = \bar{a}\sigma_3$. Then, the constraint equation (3) can be solved for $a, \bar{a}$ such that

$$\begin{aligned} a &= \frac{u^*\partial u - u\partial u^*}{4(1-uu^*)} - \frac{i}{2}\partial\theta \\ \bar{a} &= \frac{u\bar{\partial}u^* - u^*\bar{\partial}u}{4(1-uu^*)} - \frac{i}{2}\bar{\partial}\theta, \end{aligned} \tag{6}$$

which may be used to bring Eqs.(2) - (4) into a more conventional form of the complex sine-Gordon equation,

$$\begin{aligned} \partial\bar{\partial}u + \frac{u^*\partial u\bar{\partial}u}{1-uu^*} + 4\beta u(1-uu^*) &= 0 \\ \partial\bar{\partial}u^* + \frac{u\partial u^*\bar{\partial}u^*}{1-uu^*} + 4\beta u^*(1-uu^*) &= 0. \end{aligned} \tag{7}$$

After the integration over $A, \bar{A}$, the action (1) reduces to

$$\begin{aligned} I \;=\; & \frac{1}{4\pi}\int\Big[\frac{\partial u\bar{\partial}u^* + \bar{\partial}u\partial u^*}{1-uu^*} - i\partial\theta(u\bar{\partial}u^* - u^*\bar{\partial}u) - i\bar{\partial}\theta(u^*\partial u - u\partial u^*)\Big] \\ & + \frac{1}{2\pi}\int(\partial u\bar{\partial}u^* - \bar{\partial}u\partial u^*)\ln(-i\sqrt{1-uu^*}e^{-i\theta}) - \frac{\beta}{\pi}\int(2uu^*-1)\;. \end{aligned} \tag{8}$$

The second term with a logarithmic expression breaks the reality condition, whose peculiar behavior arises from the fact that it comes from the Wess-Zumino term. However, this term does not contribute to the equation of motion in Eq.(7) but is needed for the current conservation in Eq.(38). As we see below, the nonreality of the term does not cause any physical problems. The singularity at $uu^* = 1$ comes from the elimination of $A, \bar{A}$ and is not a real singularity of the theory. At $uu^* = 1$, A and $\bar{A}$ become indeterminate from the constraint equation thereby causing the aparrent sigularity over the integration. Since $uu^* = 1$ is precisely the vacuum of the theory for $\beta < 0$, it shows that the WZW actional formulation of the complex sine-Gordon theory is more natural than the conventional expression in terms of local fields u and u^* only. Moreover, we show that it reveals some salient features of the theory when we introduce the following gauge fixing; the U(1) vector gauge transform $g \rightarrow \exp(i\phi\sigma_3) g \exp(-i\phi\sigma_3)$ changes θ in Eq.(5) by $\theta + 2\phi$ while leaving u and the complex sine-Gordon equation (7) itself invariant. Owing to Eq.(4), we may fix the gauge by setting $A = \bar{A} = 0$. This is possible when the holonomy of the flat connection is trivial which is the case for the flat Minkowski base manifold. The constraint equation in the gauge $A = \bar{A} = 0$ becomes

$$\frac{u^*\partial u - u\partial u^*}{4(1-uu^*)} - \frac{i}{2}\partial\theta = 0 \ , \ \frac{u\bar{\partial} u^* - u^*\bar{\partial} u}{4(1-uu^*)} - \frac{i}{2}\bar{\partial}\theta = 0 \tag{9}$$

which may be solved for θ by integration,

$$\theta = \frac{1}{2i}\int dz \frac{u^*\partial u - u\partial u^*}{1-uu^*} + d\bar{z}\frac{u\bar{\partial} u^* - u^*\bar{\partial} u}{1-uu^*}. \tag{10}$$

In the gauge $A = \bar{A} = 0$, we assert that the theory is symmetric under the duality transform;

$$g \leftrightarrow i\sigma_1 g \ , \ \beta \leftrightarrow -\beta \tag{11}$$

which interchanges the sign of the coupling constant β. It is easy to check that the equation of motion (2) and the constraint equation (3) with $A = \bar{A} = 0$ are both invariant under the duality transform. In components, if we define the dual field g^D by

$$g^D \equiv \begin{pmatrix} v & i\sqrt{1-v\bar{v}}e^{i\phi} \\ i\sqrt{1-v\bar{v}}e^{-i\phi} & \bar{v} \end{pmatrix} = i\sigma_1 g = \begin{pmatrix} -\sqrt{1-uu^*}e^{i\theta} & iu^* \\ iu & -\sqrt{1-uu^*}e^{i\theta} \end{pmatrix}, \tag{12}$$

the duality transform is expressed by

$$v = -\sqrt{1-uu^*}e^{-i\theta}, \ u = \sqrt{1-v\bar{v}}e^{-i\phi} \text{ and } \beta \leftrightarrow -\beta \ . \tag{13}$$

The theory is also symmetric under another type of duality transform,

$$g \leftrightarrow i g\sigma_1 \ , \ \beta \leftrightarrow -\beta \tag{14}$$

which is identical with Eq.(11) except for the change of the sign of θ. The duality in Eq.(11) may be compared with the Krammers-Wannier duality of the Z_N parafermion theory bewteen the spin variables s_j and the dual spin variables μ_j. These spin variables are given by $s_k = \phi^{(k)}_{k,k}$ and $\mu_k = \phi^{(k)}_{-k,k}$ where $\phi^{(2j)}_{2m,2\bar{m}}(-j \geq m, \bar{m} \geq j \ \ ; \ j = 0, 1/2, \cdots, N/2)$ are the primary fields of the Z_N parafermion theory which is related to those of the WZW theory by

$$\Phi^{(j)}_{m,\bar{m}}(z,\bar{z}) = \phi^{(2j)}_{2m,2\bar{m}}(z,\bar{z}) : \exp\Big[\frac{im}{\sqrt{N}}\chi(z) + \frac{i\bar{m}}{\sqrt{N}}\bar{\chi}(\bar{z})\Big] : . \tag{15}$$

where $\chi(z), \bar{\chi}(\bar{z})$ are U(1) scalar fields[10][11]. In the case $j = 1$, the primary fields of WZW theory $\Phi^{(1)}$ may be given by the adjoint representation of g,

$$\Phi^{(1)}_{ab} = \mathrm{Tr} g^{-1} T_a g T_b \tag{16}$$

where T_a are generators of the algebra $su(2)$. In the limit $N \rightarrow \infty$, Eq.(15) shows that the spin variables can be written directly in terms of the WZW primary fields such that

$$\begin{aligned} s_2 &= \Phi^1_{1,1} = \mathrm{Tr} g^{-1} T_+ g T_+ = (1 - uu^*) e^{-2i\theta} \\ \mu_2 &= \Phi^1_{-1,1} = \mathrm{Tr} g^{-1} T_- g T_+ = u^2 \end{aligned} \tag{17}$$

where $T_\pm \equiv (\sigma_1 \pm i\sigma_2)/2$. Thus, the Krammers-Wannier duality which interchange s_2 and μ_2 agrees precisely with the duality transform in Eq.(13). If we assume that $u = u^* = \cos\varphi$, then the complex sine-Gordon equation reduces to the well-known sine-Gordon equation $\partial\bar{\partial}\varphi = 2\beta \sin 2\varphi$ and the duality transform reduces to the more familar one,

$$\cos\varphi \leftrightarrow \sin\varphi \ \text{ or } \ \varphi \leftrightarrow \varphi + (n + \frac{1}{2})\pi, \ n \in Z \ ; \ \beta \leftrightarrow -\beta. \tag{18}$$

The duality transform also relates exact solutions of the $\beta < 0$ case with those of the $\beta > 0$ case. To better understand the vacuum structure of the theory, we introduce another parametrization of g which is equivalent to Eq.(5),

$$g = e^{i\eta\sigma_3} e^{i\varphi(\cos\theta\sigma_1 + \sin\theta\sigma_2)} e^{i\eta\sigma_3} = \begin{pmatrix} e^{2i\eta}\cos\varphi & i\sin\varphi e^{-i\theta} \\ i\sin\varphi e^{i\theta} & e^{-2i\eta}\cos\varphi \end{pmatrix} . \tag{19}$$

The potential term then becomes

$$V = \frac{\beta}{\pi}\cos 2\varphi \tag{20}$$

so that the vacuua for $\beta < 0$ are specified by

$$\varphi = n\pi, \ n \in Z \ ; \ g_{\mathrm{vac}} = \begin{pmatrix} \pm e^{i\Omega} & 0 \\ 0 & \pm e^{-i\Omega} \end{pmatrix}, \tag{21}$$

while those for $\beta > 0$ are

$$\varphi = (n + \frac{1}{2})\pi, \ n \in Z \ ; \ g_{\mathrm{vac}} = \begin{pmatrix} 0 & \pm i e^{-i\Omega} \\ \pm i e^{i\Omega} & 0 \end{pmatrix} . \tag{22}$$

It is easy to check that these two vacuum solutions are related by the duality transform. In order to adjust the vacuum to possess zero potential energy, one has to add a constant term to the potential in which case, the coupling constant β can be treated as a mass parameter in the perturbative approach. The degeneracy of the vacuua predicts the existence of a soliton solution with a topological soliton number $\Delta n = n_1 - n_2$ which interpolates two different vacuua characterized by integers n_1 and n_2. In the following, we derive solitons explicitly from the Bäcklund transform for the complex sine-Gordon theory. The Bäcklund transform can be compactly given by

$$\Psi_g(\lambda) = \frac{\lambda}{\lambda - i\delta/\sqrt{|\beta|}}\Big(1 + \frac{\delta}{\sqrt{|\beta|}\lambda} g^{-1}\bar{T} f\Big)\Psi_f(\lambda) \ , \tag{23}$$

where δ is an arbitrary parameter and Ψ_g, Ψ_f satisfy the linear equations,

$$\begin{aligned} &(\partial + g^{-1}\partial g + \beta\lambda T)\Psi_g = 0 \quad , \quad (\bar\partial + \frac{1}{\lambda} g^{-1}\bar T g)\Psi_g = 0 \\ &(\partial + f^{-1}\partial f + \beta\lambda T)\Psi_f = 0 \quad , \quad (\bar\partial + \frac{1}{\lambda} f^{-1}\bar T f)\Psi_f = 0. \end{aligned} \tag{24}$$

If we eliminate Ψ_g, Ψ_f from Eqs.(23) and (24), we obtain the Bäcklund transform in terms of g and f only,

$$\begin{aligned} g^{-1}\partial g - f^{-1}\partial f - \frac{\delta\beta}{\sqrt{|\beta|}}[\, g^{-1}\bar T f \ , \ T \,] &= 0 \\ \bar\partial g g^{-1}\bar T - \bar T \bar\partial f f^{-1} + \frac{\sqrt{|\beta|}}{\delta}(g f^{-1}\bar T - \bar T g f^{-1}) &= 0 \ . \end{aligned} \tag{25}$$

If we apply $\bar\partial$ to the first equation in Eq.(25) and combine with the second equation, we see that g and f both satisfy the complex sine-Gordon equation. It is easy to convince that Eq.(25) is also consistent with the constraint equation (3) for $A = \bar A = 0$. Thus, the Bäcklund transform generates a new solution g from a known solution f via the first order differential equation (25). In particular, 1-soliton solution can be obtained by applying the Bäcklund transform to the vacuum solutions in Eqs.(21) and (22). For $\beta < 0$, Eq.(25), with $f = g_{\rm vac}$ in Eq.(21), becomes in components

$$\begin{aligned} \partial u + 2\delta\sqrt{-\beta}e^{i\Omega}(1 - uu^*) &= 0 \\ \bar\partial u - \frac{2\sqrt{-\beta}}{\delta}e^{i\Omega}(1 - uu^*) &= 0 \end{aligned} \tag{26}$$

and their complex conjugates. These equations and Eq.(10) may be readily integrated to yield the 1-soliton solution,

$$\begin{aligned} u &= -e^{i\Omega}\cos(\alpha)\tanh[2\sqrt{-\beta}\cos(\alpha)\frac{x - Vt}{\sqrt{1 - V^2}}] - ie^{i\Omega}\sin(\alpha) \\ \theta &= -2\sqrt{-\beta}\sin(\alpha)\frac{t - Vx}{\sqrt{1 - V^2}} \ . \end{aligned} \tag{27}$$

where $V \equiv (1-\delta^2)/(1+\delta^2)$ is the velocity of the soliton, $\cos\alpha$ and $\sin\alpha; -\pi \le \alpha < \pi$ are the constants of integration and $t \equiv \frac{1}{2}(z + \bar z), \ x \equiv \frac{1}{2}(z - \bar z)$. Similarly, we obtain the Bäcklund equation for $\beta > 0$ from the vacuum in Eq.(22),

$$\begin{aligned} \partial u - 2\delta\sqrt{\beta}e^{i(\theta+\Omega)}u\sqrt{1 - uu^*} &= 0 \\ \bar\partial u + \frac{2\sqrt{\beta}}{\delta}e^{-i(\theta+\Omega)}u\sqrt{1 - uu^*} &= 0 \ , \end{aligned} \tag{28}$$

from which we obtain the 1-soliton for $\beta > 0$,

$$\begin{aligned} u &= \frac{\cos(\alpha)}{\cosh[2\sqrt{\beta}\cos(\alpha)\frac{x-Vt}{\sqrt{1-V^2}}]}\exp(2i\sqrt{\beta}\sin(\alpha)\frac{t-Vx}{\sqrt{1-V^2}}) \\ \theta &= -\Omega - \tan^{-1}(\tan(\alpha)\coth[2\sqrt{\beta}\cos(\alpha)\frac{x-Vt}{\sqrt{1-V^2}}]) \ . \end{aligned} \tag{29}$$

This agrees with the 1-soliton for $\beta < 0$ through the duality transform. The u-part of the soliton solutions are obtained previously by Lund and Regge[1] for $\beta < 0$ and by Getmanov[12] for $\beta > 0$. Before discussing the physical meaning of the 1-soliton solutions, we note that 2-solitons and breather solutions can be also obtained from the following nonlinear superposition rule; let g_1 and g_2 be two solutions generated by the Bäcklund transform from a known solution g_0 with respect to the Bäcklund parameters δ_1 and δ_2. If we take the Bäcklund transform again on g_2 with the parameter δ_1 and also on g_1 with δ_2 and require the commutability of the process, i.e. require that both of them result in the same g_3, then we have from Eq.(23)

$$\begin{aligned} \Psi_{g_3} &= \frac{\lambda}{\lambda - i\delta_2/\sqrt{|\beta|}}\frac{\lambda}{\lambda - i\delta_1/\sqrt{|\beta|}}(1+\frac{\delta_2}{\sqrt{|\beta|}\lambda}g_3^{-1}\bar{T}g_1)(1+\frac{\delta_1}{\sqrt{|\beta|}\lambda}g_1^{-1}\bar{T}g_0)\Psi_{g_0} \\ &= \frac{\lambda}{\lambda - i\delta_1/\sqrt{|\beta|}}\frac{\lambda}{\lambda - i\delta_2/\sqrt{|\beta|}}(1+\frac{\delta_1}{\sqrt{|\beta|}\lambda}g_3^{-1}\bar{T}g_2)(1+\frac{\delta_2}{\sqrt{|\beta|}\lambda}g_2^{-1}\bar{T}g_0)\Psi_{g_0} \ . \end{aligned} \tag{30}$$

This may be solved for g_3 in terms of g_1 and g_2 such that

$$g_3 = \bar{T}(\delta_1 g_2 - \delta_2 g_1)g_0^{-1}\bar{T}^{-1}(\delta_1 g_1^{-1} - \delta_2 g_2^{-1})^{-1} \ , \tag{31}$$

which represents a nonlinear superposition rule for the complex sine-Gordon theory. In particular, if we choose g_0 as the vacuum solution in Eq.(21) then g_1 and g_2 become

$$\begin{aligned} g_k &= \begin{pmatrix} u_k & i\sqrt{1-u_k u_k^*}e^{i\theta_k} \\ i\sqrt{1-u_k u_k^*}e^{-i\theta_k} & u_k^* \end{pmatrix} \\ u_k &= -e^{i\Omega}(\cos(\alpha_k)\tanh[2\sqrt{-\beta}\cos(\alpha_k)\Sigma_k] + i\sin(\alpha_k)) \\ \theta_k &= -2\sqrt{-\beta}\sin(\alpha_k)\Theta_k \ ; \ k = 1, 2 \end{aligned} \tag{32}$$

whereas g_3 is given by Eq.(31). Σ_k and Θ_k are to be determined by the choice of parameters δ_1 and δ_2. Following three specific choices of parameters are of interest;

i) soliton-soliton scattering;

$$\delta_1 = -\frac{1}{\delta_2} = \sqrt{\frac{1-V}{1+V}}, \ \Sigma_1 = \frac{x-Vt}{\sqrt{1-V^2}}, \ \Theta_1 = \frac{t-Vx}{\sqrt{1-V^2}} \ ; \ \Sigma_2 = \frac{-x-Vt}{\sqrt{1-V^2}}, \ \Theta_2 = \frac{-t-Vx}{\sqrt{1-V^2}} \tag{33}$$

ii) soliton-antisoliton scattering

$$\delta_1 = \frac{1}{\delta_2} = \sqrt{\frac{1-V}{1+V}},\ \Sigma_1 = \frac{x-Vt}{\sqrt{1-V^2}},\ \Theta_1 = \frac{t-Vx}{\sqrt{1-V^2}}\ ;\ \Sigma_2 = \frac{x+Vt}{\sqrt{1-V^2}},\ \Theta_2 = \frac{t+Vx}{\sqrt{1-V^2}} \tag{34}$$

iii) breather

$$\delta_1 = \frac{1}{\delta_2} = \sqrt{\frac{1-iV}{1+iV}},\ \Sigma_1 = \frac{x-iVt}{\sqrt{1+V^2}},\ \Theta_1 = \frac{t-iVx}{\sqrt{1+V^2}}\ ;\ \Sigma_2 = \frac{x+iVt}{\sqrt{1+V^2}},\ \Theta_2 = \frac{t+iVx}{\sqrt{1+V^2}} \tag{35}$$

which can be shown to agree with the sine-Gordon cases when $\alpha_k = 0$.

Finally, we address the physical meaning of the parameter α which can be inferred from the global U(1) symmetries of the theory. Recall that the complex sine-Gordon theory as defined in Eq.(1) is invariant under the local vector U(1) and the global axial U(1) transformations. After the gauge fixing $A = \bar{A} = 0$, the residual symmetries are the global vector and axial U(1). The global axial vector transformation is

$$g \rightarrow e^{i\epsilon\sigma_3} g e^{i\epsilon\sigma_3} \tag{36}$$

or, in components,

$$u \rightarrow u e^{2i\epsilon}\ ,\ u^* \rightarrow u^* e^{-2i\epsilon}\ ,\ \theta \rightarrow \theta\ . \tag{37}$$

The conserved axial currents $J_A, \bar{J}_A$ satisfying the relation, $\bar{\partial} J_A + \partial \bar{J}_A = 0$, can be obtained directly from the action in Eq.(8) by the Noether method,

$$\begin{aligned} J_A &= i\frac{u^*\partial u - u\partial u^*}{1-uu^*} + 2\partial(uu^*\theta) + i\partial F(uu^*) \\ \bar{J}_A &= i\frac{u^*\bar{\partial} u - u\bar{\partial} u^*}{1-uu^*} - 2\bar{\partial}(uu^*\theta) - i\bar{\partial} F(uu^*), \end{aligned} \tag{38}$$

where the function F satisfies $F'(x) = 2\ln(-i\sqrt{1-x})$. We caution that the action in Eq.(8) is an effective action where the constraint Eq.(3) is imposed. The Noether method can be applied consistently only when the constraint is invariant under the axial transformation which is indeed the case. The conserved axial charge arising from currents in Eq.(38) is

$$Q_A = -i\int_{-\infty}^{\infty} dx \frac{u^*\partial_t u - u\partial_t u^*}{1-uu^*} - \Big[2uu^*\theta + iF(uu^*)\Big]_{-\infty}^{\infty}\ . \tag{39}$$

Thus the axial charges of 1-soliton solutions computed in the rest frame are

$$Q_A(\beta < 0) = 0\ ;\ Q_A(\beta > 0) = 4(\text{sign}[\alpha]\frac{\pi}{2} - \alpha)\ \text{for}\ \sin\alpha \neq 0 \tag{40}$$

which shows that the parameter α characterizes the U(1) charge of solitons. If $\sin\alpha = 0$, the axial charge is indeterminate because of the singular behavior of Q_A at $uu^* = 1$. As stated earlier, this relects the fact that $A, \bar{A}$ are indeterminate at $uu^* = 1$. Here, we simply take Q_A to be zero. This may be done more rigorously by taking an appropriate limiting procedure in the gauge $A = \bar{A} = 0$, which however is physically irrelevant. In the case of the vector U(1) symmetry, we note that the duality transform in Eq.(11) relates the axial symmetry with the vector symmetry in the following way; if we perform the axial transform in Eq.(36), then the dual field $g^D = i\sigma_1 g$ changes according to the vector transform,

$$g^D \to e^{-i\epsilon\sigma_3} g^D e^{i\epsilon\sigma_3} \ . \tag{41}$$

This implies that the conserved vector currents and charges are dual to the axial ones. That is, the vector charges Q_V are given by

$$Q_V(\beta < 0) = Q_A(\beta > 0) \ ; \ Q_V(\beta > 0) = Q_A(\beta < 0) \tag{42}$$

whereas the vector currents $J_V, \bar{J}_V$ can be obtained by taking the dual transform of the axial currents such that

$$\begin{aligned} J_V &= \partial(iuu^* \ln \frac{u^*}{u} + iF) \\ \bar{J}_V &= -\bar{\partial}(iuu^* \ln \frac{u^*}{u} + iF) \ . \end{aligned} \tag{43}$$

Another aspect of the parameter α is that it dictates the topological nature of solitons. If $\cos\alpha = 0$, the solutions in Eqs.(27) and (29) trivially reduce to the vacuum solutions in Eqs.(21) and (22). If $\sin\alpha = 0$, it is easy to check that the analyticity of the solution requires that $\Delta n = \pm 1$, i.e. it becomes a topological 1-soliton (antisoliton). For $\cos\alpha \neq 0$ and $\sin\alpha \neq 0$, the continuity of the solution requires that $\Delta n = 0$, which shows that charged solitons are nontopological. Nevertheless, they are stable against the decay into multi-nontopological solitons and mesons. This semiclassical stability may be shown by the mass formula[4] for the nontopological soliton of charge Q[8]

$$M(Q) = \frac{4m}{\lambda^2} |\sin \frac{\lambda^2 Q}{4}|, \ |Q| \leq \frac{2\pi}{\lambda^2} \ . \tag{44}$$

It is easy to check that $M(Q_1) + M(Q_2) > M(Q_1 + Q_2)$ which prohibits the nontopological soliton from decaying in two solitons conserving the charge. Also, the decay into the elementary meson of charge 1 and mass m is suppressed by the bound $M(1) = \frac{4m}{\lambda^2} \sin \frac{\lambda^2}{4} < m$.

ACKNOWLEDGEMENT

We would like to thank I.Bakas, R. Sasaki and T.Hollowood for discussion. This work was supported in part by the program of Basic Science Research, Ministry of Education BSRI-95, and by Korea Science and Engineering Foundation through CTP/SNU and the Korea-Japan Cooperative Science Program.

---

[4] Here, the mass constant m is our coupling constant $2\sqrt{|\beta|}$. The constant λ^2 gets renormalized in the quantum case which in our classical consideration has been set to one.

References

[1] F.Lund and T.Regge, Phys.Rev. **D14** 1524 (1976).

[2] K.Pohlmeyer, Commun.Math.Phys.**46** 207 (1976).

[3] I.Bakas, Int.J.Mod.Phys. **A 9** (1994) 3443.

[4] Q-H.Park, Phys.Lett.**B328** 329 (1994).

[5] Q-H.Park and H.J.Shin, Phys.Lett.**B347** 73 (1995).

[6] T.J.Hollowood, J.L.Miramontes and Q-H.Park, "Massive integrable soliton theories", to appear in Nucl.Phys.B.

[7] Q-Han Park and H.J.Shin, "Classical matrix sine-Gordon theory", (Kyung Hee U.) SNUCTP-95-55, hep-th/9505017.

[8] N. Dorey, T.J. Hollowood "Quantum scattering of charged solitons in the complex sine-Gordon model" (Swansea U.). SWAT-46, hep-th/9410140.

[9] A. Zamolodchikov and V. Fateev, Sov.Phys.JETP **62** (1985) 215; Sov.J.Nucl.Phys. **43** (1986) 657.

[10] V. Knizhnik and A. Zamolodchikov, Nucl.Phys.**B247** (1984) 83.

[11] D. Gepner and E. Witten, Nucl.Phys.**B278** (1986) 493.

[12] B. Getmanov, Sov.Phys. JETP Lett. **25** (1977) 119.